

Non-Abelian Class Field Theory for Riemann Surfaces

Lin WENG

Graduate School of Mathematics, Nagoya University, Japan

In this paper, using what we call a micro reciprocity law, we complete Weil's program [W] for non-abelian class field theory of Riemann surfaces.

1. Refined Structures for Tannakian Categories

Let $\mathbf{T}$ be a Tannakian category with a fiber functor $\omega : \mathbf{T} \rightarrow \text{Ver}_{\mathbf{C}}$, where $\text{Ver}_{\mathbf{C}}$ denotes the category of finite dimensional $\mathbf{C}$ -vector spaces. An object $t \in \mathbf{T}$ is called *reducible* if there exist non-zero objects $x, y \in \mathbf{T}$ such that $t = x \oplus y$. An object is called *irreducible* if it is not reducible. If moreover every object x of $\mathbf{T}$ can be written uniquely as a sum of irreducible objects $x = x_1 \oplus x_2 \oplus \dots \oplus x_n$, then $\mathbf{T}$ is called a *unique factorization* Tannakian category. Usually, we call x_i 's the *irreducible components* of x .

A Tannakian subcategory $\mathbf{S}$ of a unique factorization Tannakian category $\mathbf{T}$ is called *completed* if for $x \in \mathbf{S}$, all its irreducible components x_i 's in $\mathbf{T}$ are also in $\mathbf{S}$. $\mathbf{S}$ is called *finitely generated* if as a Tannakian category, it is generated by finitely many objects. Moreover, $\mathbf{S}$ is called *finitely completed* if (a) $\mathbf{S}$ is finitely generated; (b) $\mathbf{S}$ is completed; and (c) $\text{Aut}^{\otimes} \omega|_{\mathbf{S}}$ is a finite group.

2. An Example

Let M be a compact Riemann surface of genus g . Fix an effective divisor $D = \sum_{i=1}^N e_i P_i$ with $e_i \in \mathbf{Z}_{\geq 2}$ once for all. For simplicity, in this note we always assume that $(M; D) \neq (\mathbf{P}^1; e_1 P_1)$, or $(\mathbf{P}^1; e_1 P_1 + e_2 P_2)$ with $e_1 \neq e_2$. (These cases may be easily treated.)

By definition a parabolic semi-stable bundle

$$\Sigma := (E =: E(\Sigma); P_1, \dots, P_N; \text{Fil}(E|_{P_1}), \dots, \text{Fil}(E|_{P_N}); a_{11}, \dots, a_{1r_1}; \dots; a_{N1}, \dots, a_{Nr_N})$$

of parabolic degree 0 is called a *GA bundle* over M along D if (i) the parabolic weights are all rational, i.e., $a_{ij} \in \mathbf{Q} \cap [0, 1)$; (ii) there exist $\alpha_{ij} \in \mathbf{Z}, \beta_{ij} \in \mathbf{Z}_{>0}$ such that (a) $(\alpha_{ij}, \beta_{ij}) = 1$; (b) $a_{ij} = \alpha_{ij}/\beta_{ij}$; and (c) $\beta_{ij}|e_i$, for all i, j . Denote by $[\Sigma]$ the Seshadri equivalence class associated with Σ . Moreover, for $[\Sigma]$, define $\omega_D([\Sigma])$ as $E(\text{Gr}(\Sigma))|_P$, i.e., the fiber of the bundles associated with Jordan-Hölder graded parabolic bundle of Σ at a fixed $P \in M^0 := M \setminus |D| = M \setminus \{P_1, \dots, P_N\}$.

Proposition. *With the same notation as above, put $\mathcal{M}(M; D) := \{[\Sigma] : \Sigma \text{ is a GA bundle over } M \text{ along } D\}$. Then $\mathcal{M}(M; D)$ is a unique factorization Tannakian category and $\omega_D : \mathcal{M}(M; D) \rightarrow \text{Vec}_{\mathbf{C}}$ is a fiber functor.*

Proof. (1) By a result of Mehta-Seshadri [MS, Prop. 1.15], $\mathcal{M}(M; D)$ is an abelian category. Then from the unitary representation interpretation of a GA bundle, a fundamental result due to Seshadri, (see [MS, Thm 4.1], also in Step 3 of Section 5 below,) $\mathcal{M}(M; D)$ is closed under tensor product. The rigidity may be checked directly.

(2) Since the Jordan-Hölder graded bundle is a direct sum of stable and hence irreducible objects and is unique, (see [MS, Rm 1.16],) so, $\mathcal{M}(M; D)$ is a unique factorization category.

(3) By definition, we know that the functor ω is exact and tensor. So we should check whether it is faithful. This then is a direct consequence of the fact that $\mathcal{M}(M; D)$ is a unique factorization category and that any morphism between two irreducible objects is either zero or a constant multiple of the identity map.

3. Reciprocity Map

In $[\Sigma]$, choose its associated Jordan-Hölder graded bundle $\text{Gr}(\Sigma)$ as a representative. Then by the above mentioned fundamental result of Seshadri, $\text{Gr}(\Sigma)$ corresponds to a unitary representation $\rho_{\text{Gr}(\Sigma)} : \pi_1(M^0) \rightarrow U(r_{\Sigma})$, where r_{Σ} denotes the rank of $E(\Sigma) := E$.

For each element $g \in \pi_1(M^0)$, we then obtain a $\mathbf{C}$ -isomorphism of $E(\text{Gr}(\Sigma))|_P$. Thus, in particular, we get a natural morphism

$$W : \pi_1(M^0) \rightarrow \text{Aut}^{\otimes} \omega_D.$$

Now note that $\pi_1(M^0)$ is generated by $2g$ hyperbolic transformations $A_1, B_1, \dots, A_g, B_g$ and N parabolic transformations $S_1, \dots, S_N$ satisfying a single relation $A_1 B_1 A_1^{-1} B_1^{-1} \dots A_g B_g A_g^{-1} B_g^{-1} S_1 \dots S_N = 1$, and that $\rho_{\text{Gr}(\Sigma)}(S_i^{e_i}) = 1$ for all $i = 1, \dots, N$. ([MS, §1].) Denote by $J(D)$ the normal subgroup of $\pi_1(M^0)$ generated by $S_1^{e_1}, \dots, S_N^{e_N}$. Then naturally we obtain the following *reciprocity map*

$$W(D) : \pi_1(M^0)/J(D) \rightarrow \text{Aut}^{\otimes} \omega_D.$$

4. Main Theorem

As usual, a Galois covering $\pi : M' \rightarrow M$ is called branched at most at D if (1) π is branched at $P_1, \dots, P_N$; and (2) the ramification index e'_i of points over P_i divides e_i for all $i = 1, \dots, N$. Clearly, by changing D , we get all finite Galois coverings of M .

Main Theorem. (1) (Existence and Conductor Theorem) *There is a natural one-to-one correspondence w_D between*

$$\{\mathbf{S} : \text{finitely completed Tannakian subcategory of } \mathcal{M}(M; D)\}$$

and

$$\{\pi : M' \rightarrow M : \text{finite Galois covering branched at most at } D\};$$

(2) (Reciprocity Law) *There is a natural group isomorphism*

$$\text{Aut}^{\otimes}(\omega_D|_{\mathbf{S}}) \simeq \text{Gal}(w_D(\mathbf{S})).$$

5. Proof

Step 1: Galois Theory. By a result of Bongaard, Nielsen, Fox, M. Kato, and Namba, (see e.g., [Na, Thms 1.2.15 and 1.3.9].) we know that the assignment $(\pi : M' \rightarrow M) \mapsto \pi_*(\pi_1(M' \setminus \pi^{-1}\{P_1, \dots, P_N\}))$ gives a one-to-one correspondence between isomorphism classes of finite Galois coverings $\pi : M' \rightarrow M$ branched at most at D and finite index (closed) normal subgroups $K = K(\pi)$ of $\pi_1(M^0)$ containing $J(D)$. Moreover, we have a natural isomorphism $\text{Gal}(\pi) \simeq \pi_1(M^0)/K(\pi) \left(\simeq (\pi_1(M^0)/J(D)) / (K(\pi)/J(D)) \right)$. Thus the problem is transformed to the one for finite index normal subgroups of $\pi_1(M^0)$ which contain $J(D)$, or the same, finite index normal subgroups of $G(D) := \pi_1(M^0)/J(D)$.

Step 2: Namba Correspondence. Consider now the category $\mathbf{T}(D)$ of equivalence classes of unitary representations of $G(D)$. Clearly, $\mathbf{T}(D)$ forms a unique factorization Tannakian category, whose fiber functor $\omega(D)$ may be defined to be the forget functor. Now fixed once for all a representative $\rho_{\Sigma} : G(D) \rightarrow U(r_{\Sigma})$ for each equivalence classes $[\Sigma]$. (The choice of the representative will not change the essentials below as the resulting groups are isomorphic to each other.)

Let $\mathbf{S}$ be a finitely completed Tannakian subcategory of $\mathbf{T}(D)$. Then as in the definition of reciprocity map above, we have a natural morphism $G(D) \xrightarrow{\omega_{\mathbf{S}}} \text{Aut}^{\otimes} \omega|_{\mathbf{S}}$. Denote its kernel by $K(\mathbf{S})$. Then, by definition, $G(D)/K(\mathbf{S})$ is a finite group, and $K(\mathbf{S}) = \bigcap_{[\Sigma] \in \mathbf{S}} \ker \rho_{\Sigma}$.

Since $\mathbf{S}$ is finitely completed, there exists a finite set $S = \{[\Sigma_1], [\Sigma_2], \dots, [\Sigma_t]\}$ which generates $\mathbf{S}$ as a completed Tannakian subcategory. Set $[\Sigma_0] := \bigoplus_{i=1}^t [\Sigma_i]$. Then for any $[\Sigma] \in \mathbf{S}$, $\ker(\rho_{\Sigma_0}) \subset \ker(\rho_{[\Sigma]})$, since $\mathbf{S}$ is generated by S . Also, by definition, $\ker(\rho_{\Sigma_0}) = \bigcap_{i=1}^t \ker(\rho_{\Sigma_i})$. Thus $K(\mathbf{S}) = \ker(\rho_{\Sigma_0})$.

Therefore, for any $[\Sigma] \in \mathbf{S}$, $\rho_{\Sigma} = \tilde{\rho}_{\Sigma} \circ \Pi(D; \mathbf{S})$ where $\Pi(D; \mathbf{S}) : G(D) \rightarrow G(D)/K(\mathbf{S})$ denotes the natural quotient map and $\tilde{\rho}_{\Sigma}$ is a suitable unitary representation of $G(D)/K(\mathbf{S})$.

Now set $\tilde{\mathbf{S}} := \{[\tilde{\rho}_{\Sigma}] : [\Sigma] \in \mathbf{S}\}$. $\tilde{\mathbf{S}}$ is a finitely completed Tannakian subcategory in $\text{Uni}(G(D)/K(\mathbf{S}))$, the category of equivalence classes of unitary representations of $G(D)/K(\mathbf{S})$.

In particular, since the unitary representation $\tilde{\rho}_{\Sigma_0}$ of $G(D)/K(\mathbf{S})$ maps $G(D)/K(\mathbf{S})$ injectively into its image, for any two elements $g_1, g_2 \in G(D)/K(\mathbf{S})$, $\tilde{\rho}_{\Sigma_0}(g_1) \neq \tilde{\rho}_{\Sigma_0}(g_2)$.

With this, by applying the van Kampen Completeness Theorem ([Ka]), which claims that for any compact group G , if Z is a subset of the category $\text{Uni}(G)$ such that for any two elements g_1, g_2 , there exists a representation ρ_{g_1, g_2} in Z such that $\rho_{g_1, g_2}(g_1) \neq \rho_{g_1, g_2}(g_2)$, then the completed Tannakian subcategory

generated by Z is the whole category $\mathcal{U}ni(G)$ itself, we conclude that $\tilde{\mathbf{S}} = \mathcal{U}ni(G(D)/K(\mathbf{S}))$. But as categories, $\mathbf{S}$ is equivalent to $\tilde{\mathbf{S}}$, thus by the Tannaka duality, (see e.g., [DM] or [Ta]) we obtain a natural isomorphism $\text{Aut}^{\otimes} \omega|_{\mathbf{S}} \simeq G(D)/K(\mathbf{S})$.

On the other hand, if K is a finite index (closed) normal subgroup of $G(D)$. Set $\tilde{\mathbf{S}} := \mathcal{U}ni(G(D)/K)$ with the fiber functor $\omega_{\tilde{\mathbf{S}}}$. Compositing with the natural quotient map $\Pi : G(D) \rightarrow G(D)/K$ we then obtain an equivalent category $\mathbf{S}$ consisting of corresponding unitary representations of $G(D)$. $\mathbf{S}$ may also be viewed as a Tannakian subcategory of $\mathcal{U}ni(G(D))$. We next show that indeed such an $\mathbf{S}$ is a finitely completed Tannakian subcategory.

From definition, $\text{Aut}^{\otimes} \omega|_{\mathbf{S}} \simeq \text{Aut}^{\otimes} \omega_{\tilde{\mathbf{S}}}$ which by the Tannaka duality theorem is isomorphic to $G(D)/K$. So it then suffices to show that $\mathbf{S}$ is finitely generated. But this is then a direct consequence of the fact that for any finite group there always exists a unitary representation such that the group is injectively mapped into the unitary group.

Step 3: A Micro Reciprocity Law. With Steps 1 and 2, the proof of the Main Theorem is then completed by the following

Weil-Narasimhan-Seshadri Correspondence (Seshadri, [MS, Thm 4.1]) *There is a natural one-to-one correspondence between isomorphism classes of unitary representations of fundamental groups of M^0 and equivalence classes of semi-stable parabolic bundles over M^0 of parabolic degree zero.*

Indeed, with this theorem, the Seshadri equivalence classes of GA bundles over M along D correspond naturally in one-to-one to the equivalence classes of unitary representations of the group $\pi_1(M^0)/J(D)$. Thus by Step 2, the finitely completed Tannakian subcategories of $\mathcal{M}(M; D)$ are in one-to-one correspond to the finite index closed normal subgroup of $\pi_1(M^0)/J(D)$, which by Step 1 in one-to-one correspond to the finite Galois coverings of M branched at most at D . This gives the existence theorem. Along the same line, we have the reciprocity law as well.

REFERENCES

- [DM] P. Deligne & J.S. Milne, Tannakian categories, in *Hodge Cycles, Motives and Shimura Varieties*, LNM **900**, (1982), 101-228
- [Ka] E. van Kampen, Almost periodic functions and compact groups, *Ann. of Math.* **37** (1936), 78-91
- [MS] V.B. Mehta & C.S. Seshadri, Moduli of vector bundles on curves with parabolic structures. *Math. Ann.* **248** (1980), no. 3, 205-239.
- [MFK] D. Mumford, J. Fogarty & F. Kirwan, *Geometric Invariant Theory*, Springer-Verlag, 1994
- [Na] M. Namba, *Branched coverings and algebraic functions*. Pitman Research Notes in Mathematics Series **161**, Longman Scientific & Technical, 1987
- [NS] M.S. Narasimhan & C.S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math.* **82** (1965), 540-567
- [Se1] C.S. Seshadri, Moduli of π -vector bundles over an algebraic curve. *Questions on Algebraic Varieties*, C.I.M.E., III, (1969) 139-260
- [Se2] C. S. Seshadri, *Fibrés vectoriels sur les courbes algébriques*, *Asterisque* **96**, 1982
- [Ta] T. Tannaka, *Theory of topological groups*, Iwanami, 1949 (in Japanese)
- [W] A. Weil, Généralisation des fonctions abéliennes, *J. Math Pures et Appl.* **17**, (1938) 47-87
- [We] L. Weng, A Program for Gromatric Arithmetic, preprint, 2001